



International Journal of Veterinary Science & Technology

Research article

Effect of Inclusion Levels of *Arachis glabrata* in the Diets on the Ingestion and *In Vivo* Digestibility of *Panicum maximum* in Guinea Pigs (*Cavia porcellus*) - ②

Emile Miegoue^{1*}, Fernand Tendonkeng¹, Nathalie Mweugang Ngouopo², Jules
Lemoufouet¹, Martin Vidal Tatang¹, Paulette Ntsafack¹, Mouchili Mama¹ and Etienne
Tendonkeng Pamo¹

¹University of Dschang, Faculty of Agronomy and Agricultural Sciences, Department of Animal Production
Animal Nutrition and production Research Unit B.P. 222 Dschang, Cameroon

²University of Ngaoundere, Faculty of Sciences, Department of Animal Science

***Address for Correspondence:** Emile Miegoue, University of Dschang, Faculty of Agronomy and Agricultural
Sciences, Department of Animal Production Animal Nutrition and production Research Unit B.P. 222 Dschang,
Cameroon, Tel: +237-697-121-2 97 / 674 -218-1 74; E-mail: migoumile@yahoo.fr; emile.miegoue@univ-dschang.org

Submitted: 09 November 2018; **Approved:** 28 November 2018; **Published:** 01 December 2018

Cite this article: Miegoue E, Tendonkeng F, Ngouopo NM, Lemoufouet J, Tatang MV, et al. Effect of inclusion
levels of *arachis glabrata* in the diets on the ingestion and in vivo digestibility of *panicum maximum* in guinea
pigs (*cavia porcellus*). Int J Vet Sci Technol. 2018;2(2): 026-032.

Copyright: © 2018 Miegoue E, et al. This is an open access article distributed under the Creative
Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any
medium, provided the original work is properly cited.

Abstract

This study, which evaluated the effect of graded levels of *Arachis glabrata* on the ingesting and *in vivo* digestibility of *Panicum maximum* and on the caecal rate of bacterial flora, was conducted at the Research and Application Farm of the University of Dschang between July and August 2016. For this trial, 40 adult guinea pigs of local breed including 20 animals per sex, aged about 04 months and weighing between 288 and 374 g, were purchased from breeders in the city of Dschang and its surroundings. Guinea pigs were first acclimatized for 2 weeks in the farm's dressing boxes before being randomly distributed in the individual digestibility cages. After 10 days of adaptation, animals were distributed in 04 batches (T0, T1, T2 and T3) of 10 animals (5 males and 5 females) each then subjected to 04 iso-nitrogenous rations (16% of proteins) containing graded levels of *A. glabrata*. Animals in each batch received 250g of *Panicum maximum* associated with 60g of compound food containing 0, 10, 15 or 20% of *A. glabrata* respectively for lots T0, T1, T2 and T3. During the 07day of digestibility test, each animal's feces were collected in labeled envelopes and weighed every morning before a re-distribution of diet. *P. Maximum* and nutrients ingestion (160.3 g DM / day / animal) in males and regardless of sex was significantly ($P < 0.05$) higher with diet containing *A. glabrata* compared to control diet. In contrast, ingestion of concentrate containing legume was significantly ($P < 0.05$) lower than that of the control diet, regardless of sex. However, females had significantly ($P < 0.05$) better ingested DM and OM of the control diet than males whose ingestion of these nutrients with other diets was better ($P < 0.05$). In addition, intake of CP and CF from all rations did not significantly ($P > 0.05$) varied between the two sexes. Nutrient digestion was not significantly influenced ($P > 0.05$) by sex and as inclusion level of *A. glabrata* in the diet. The level of bacteria in the caecal flora after the digestibility test was not modified ($P > 0.05$) by the inclusion level of *A. glabrata* in the diet. Nevertheless, the diet containing 20% of *A. glabrata* was the most favorable for the development of Lactobacilli at the expense of Enterobacteria.

Keywords: *Panicum maximum*; *Arachis glabrata*; Guinea pigs; Ingestion; *In vivo* digestibility

INTRODUCTION

Food security is a real challenge in most African countries where people suffer from protein malnutrition [1]. Emerging challenges such as climate change, environmental sustainability and rapidly evolving technologies are transforming food systems and raising questions about how to sustainably meet the food needs of a growing world population [2]. Average animal protein consumption in Africa is less than one quarter of that in the Americas, Europe and Oceania, and represents only 17% of the recommended level of consumption for all proteins [3]. The same author reported that livestock and livestock products can make an essential contribution to the economic and food security of low-income households and their nutrition. According to Noubissi et al. [4], setting up short-cycle farms like caviaculture proves to be one of the durable solutions, given the major interest of the guinea pig living in its prolificacy, its high growth rate and its lean meat rich in protein.

However, food is the main limiting factor for the expression of the production potential of animals in tropical environments [5]. Indeed, the high price of compound food used as a protein supplement, in addition to raising the cost of production, does not always make it possible to obtain good productivity [6]. However, to minimize feed costs for farmed guinea pigs, suggested maximizing feed intake and diet concentrates of vitamin and vitamin supplements (CMV) [7].

In this logic, several studies have been conducted on the use of *Arachis glabrata* in the guinea pig diet. Indeed, the work of Miegoue et al. [8] and Miegoue et al. [9] confirmed the good use of this legume in ingesting and *in vivo* digestibility of *Pennisetum purpureum* or *Panicum maximum* in guinea pigs. These authors have reported that an adequate association of local forages (grasses and legumes) can constitute an undeniable food base for the growth of domestic guinea pigs with regard to their strictly herbivorous diet. However, supplementation makes it possible to compensate for deficits in nutritional principles that the simple increase in the level of forage intake cannot compensate [10].

In addition, the digestibility of the wall of plant matter is highly dependent on the action of cecal microorganisms [11], which can digest fibrous foods in the same way as polygastrics [7].

However, very little work has been done on an assessment of the level of inclusion of *Arachis glabrata* on grass digestibility. In order

to contribute to a better use of *Arachis glabrata* in the guinea pig diet, the present work consisted in evaluating the effect of different levels of *Arachis glabrata* in the diet on the ingestion and digestibility of *Panicum maximum* and bacterial flora of cecum in guinea pigs.

MATERIAL AND METHODS

The study took place between August and September 2016 at the Research and Application Farm (RAF) of University of Dschang. The locality is located between 5° 25' and 5° 30' North Latitude, 10° 0' and 10° 05' East Longitude and at an altitude of about 1420 m west of Cameroon. The climate of the region is equatorial of Cameroonian type, with an average annual temperature of 20°C. The months of July and August are the coldest. The average annual rainfall varies between 1500 and 2000 mm, with a relative humidity ranging between 40% (in the dry season) and 97% (during the heavy rains). The dry season alternates with the rainy season [12].

Housing and animal sample

The walls and floor of the breeding lodges as well as the breeding equipment (feeders, drinking troughs, and digestibility cages) were previously cleaned and disinfected with a solution of sodium hypochlorite and potassium permanganate. The realization of the crawl space for two weeks preceded the introduction of guinea pigs in the breeding unit reserved for this purpose. Forty (40) local adult guinea pigs including 20 females and 20 males of mean age 04 months and mean weight 331 ± 42.86 g were used for this test. These animals were purchased from stockbreeders in the town of Dschang and surrounding areas. The guinea pigs were acclimatized on the farm for two (02) weeks in breeding lodges, males isolated from females. Guinea pigs were raised on the ground on untreated dry wood chip litter and renewed every 2 days to prevent the accumulation of feces and urine. Each lodge was equipped with a lighting device and 2 concrete drinkers in which the drinking water enriched with vitamin C at a rate of 250 mg tablet for 1.5 liters, was available ad libitum and renewed daily. It is after this period of acclimatization that animals were introduced into individual cages of digestibility to be subjected to different tests.

Plant sample

The grass (*Panicum maximum*) was harvested before flowering in the forage plot of the farm the day before and pre-washed before

being served the next day to the animals. The leaves of the legume (*Arachis glabrata*) were harvested before flowering, dried and then crushed and incorporated into the feed. A 100 g sample of each plant dried at 60°C in an oven to a constant weight was milled and stored in plastic bags for chemical composition evaluation [13] as is shown in table 2.

Conduct of the test

Formulation of different diet: The combination of the ingredients, the proportions of which are presented in table 1, made it possible to formulate four (04) iso-nitrogen diets (16% CP).

Evaluation of the ingestion: The present digestibility test was preceded by a 10-day period of animal adaptation to the digestibility cage and experimental rations. During this period, the amount of feed was adjusted to the estimated daily feed consumption of 60 g and the animals, randomly distributed in individual cages, continued to receive vitamin C in the drinking water renewed daily.

During the trial, each treatment was repeated over ten (10) guinea pigs, five (05) of each sex.

For evaluation of ingestion, 250 grams of *Panicum maximum* associated with 60 grams of compound food (CA0, CA10, CA15 or CA20) were served once a day every day between 8 and 9 hours per animal and refusals were collected and weighed before any new distribution.

Assessment of digestibility

The digestibility test was preceded by a 10-day period of animal adaptation to the digestibility cage and experimental diet. To evaluate the digestibility that lasted seven (07) days, feces were collected and weighed daily in the morning before any further distribution of the diet and a sample of one hundred (100) grams of feces was then collected and dried at 60°C in an oven to constant weight. Subsequently, the dried feces were crushed using a home-made mill and stored in plastic bags for evaluation of their Dry Matter (DM), Organic Matter (OM), Crude Protein (CP) and Crude Cellulose (CF) according to the method described by AOAC [13].

Table 1: Formulation of different rations.

Ingredients (%)	Differents Diets			
	CA0	CA10	CA15	CA20
Middlings	31	27.5	26.5	25
Corn	30	27	25.5	24
Cotton seed cake	5	4.5	4	4
Palm kernel cake	25	22.5	21	20
Soybean meal	2	2	2	1.5
Fish meal	3	2.5	2.5	2.5
Shell powder	2	2	1.5	1.5
Prémix	1	1	1	1
Cooking salt	1	1	1	1
<i>Arachis glabrata</i>	0	10	15	20
TOTAL	100	100	100	100

Table 2: Chemical composition of forage analyzed.

Chemical composition	Forages	
	<i>P. maximum</i>	<i>A. glabrata</i>
Dry matter (%)	91.76	95.29
Organic matter (%DM)	85.88	87.82
Crude protein (% DS)	13.45	19.53
Crude fiber (% DS)	33.08	20.94
Ash (% DS)	14.12	12.18

Analysis of caecal flora

At the end of the digestibility test, 03 randomly selected guinea pigs of each sex and treatment were sacrificed for evaluation of the composition of the caecal flora (Enterobacteria and Lactobacilli) according to the method described [14].

Statistical analyzes

Data on dietary intake and nutrient digestibility, as well as on the caecal flora were subjected to the 2-factor analysis of the variance (food ration and sex of the animal) according to the General Linear Model (MLG).

When significant differences existed between treatments, the separation of means was done by the Waller Duncan test at the 5% threshold [15]. The statistical analysis software used was SPSS 20.0. The comparison between the sexes was made by Student's "t" test at the 5% threshold.

RESULTS

Chemical composition of fodder and different diets

The chemical composition of these forages revealed that the dry matter, organic matter and crude protein contents of *Arachis glabrata* were higher than those of *Panicum maximum* (Table 2). On the other hand, the crude fiber and ash content of the legume was rather low compared to that of the grass.

The table 3 shows dry matter, organic matter, crude protein, digestible energy and ash contents of different diet. It showed very little variation between diets. In contrast, the fat content of the control was higher (8.74% DM) than the mean value of that obtained with diets containing *Arachis glabrata* (4.58% DM). The crude fiber content of the legume-free diet was lower than the values obtained with other diets.

Effect of inclusion level of *Arachis glabrata* in the diet on ingestion of *Panicum maximum* in guinea pigs

In male guinea pigs and regardless of sex (Table 4), ingestion of *Panicum maximum*, compound feed, total Dry Matter (DM), Organic Matter (OM), Crude Protein (CP) and Crude Fiber (CF) did not vary ($p > 0.05$) among *A. glabrata*-containing diets but significantly ($p < 0.05$) higher than that of the control diet. Compared with the legume-free diet, intake of the diet consisting of *A. glabrata*-containing diets was significantly ($p < 0.05$) lower than that of the control diet. The highest intake of DM (179.5 g DM/ animal/ day), OM (154.3 g DM/ animal/ day), CP (24.7 g DM/ animal/ day) and CF (56.1 g DM/ animal/ day) was obtained with the diet containing 20% *A. glabrata*.

Effect of sex on nutrient ingestion in guinea pigs fed with *P. maximum* associated with diets containing different levels of *A. glabrata*

It is apparent from figure 1 that sex significantly influenced ($p < 0.05$) ingestion of DM and OM in favor of females with the legume-free diet, whereas with diet containing *A. glabrata*, males are those who had better ingested the food. In contrast, CP and CF of all diets were ingested in the same manner ($p > 0.05$) between males and females.

Effect of inclusion level of *Arachis glabrata* in the diet on Apparent Digestibility Coefficients (ADC) of nutrients in guinea pigs fed with *P. maximum*

The inclusion level of *Arachis glabrata* in the diet had no significant effect ($p > 0.05$) on digestive utilization of nutrients in

males or regardless of the sex in guinea pigs fed with *P. maximum* (Table 5). It was with the control diet that the highest ADC values of DM (92.6 g DM/ animal/ day), OM (92.8 g DM/ animal/ day), CP (95.6 g DM/ animal/ j) and CF (91.1 g DM/ animal/ day) were recorded.

Effect of sex on apparent digestibility coefficients of nutrients in guinea pigs fed *P. maximum* associated with diets containing different levels of *Arachis glabrata*

Apparent Digestibility Coefficients (ADC) of nutrients were not significantly influenced ($p > 0.05$) by sex when guinea pigs were fed with *P. maximum* regardless of diet (Figure 2).

Effect of the level of inclusion of *Arachis glabrata* in diet on the caecal rate of Enterobacteria or Lactobacillus in

Table 3: analyzed chemical composition of different diets.

Chemical composition	Differents Diets			
	CA0	CA10	CA15	CA20
Dry matter (%)	97.82	97.28	97.68	97.54
Organic matter (%DM)	86.06	87.76	88.44	87.45
Crude protein (% DM)	16.02	16.43	16.55	16.67
Fat (% DM)	08.74	04.80	04.65	04.30
Crude fiber (% DM)	15.80	17.93	17.80	17.46
Ash (% DM)	13.94	12.24	11.56	12.55
Digestible energy (kcal/kg ingested food)	2690	2640	2625	2595

Table 4: Ingestion of *P. maximum* in Guinea Pigs associated with inclusion Levels of *Arachis glabrata* in the Diet.

Ingestions (gDM/d/ animal)	Treatments				ESM	P
	PMCA0	PMCA10	PMCA15	PMCA20		
Expérimental food						
Panicum maximum(PM)						
♂(5)	126.1 ^b	158.3 ^a	154.9 ^a	161.6 ^a	3.99	0.04
♀(5)	145.6 ^a	151.3 ^a	140.5 ^a	159.0 ^a	3.35	0.28
♂♀(10)	135.8 ^b	154.8 ^a	147.7 ^a	160.3 ^a	3.09	0.03
Compound food						
♂(5)	24.6 ^a	17.1 ^b	18.1 ^b	21.6 ^a	0.88	0.01
♀(5)	21.0 ^a	16.7 ^b	16.4 ^b	12.8 ^b	0.95	0.05
♂♀(10)	22.8 ^a	17.0 ^b	17.3 ^b	17.2 ^b	0.86	0.04
Nutrients						
Total dry matter (DM)						
♂(5)	150.7 ^b	178.8 ^a	173.0 ^a	183.0 ^a	3.79	0.04
♀(5)	166.7 ^a	167.8 ^a	157.0 ^a	176.1 ^a	3.57	0.36
♂♀(10)	158.7 ^b	177.3 ^a	165.0 ^a	179.5 ^a	3.04	0.03
Organic matter (OM)						
♂(5)	129.5 ^b	153.8 ^a	149.0 ^a	157.5 ^a	3.26	0.04
♀(5)	143.2 ^a	145.3 ^a	135.2 ^a	151.1 ^a	3.07	0.40
♂♀(10)	136.3 ^b	152.0 ^a	142.1 ^a	154.3 ^a	2.62	0.03
Crude protein(CP)						
♂(5)	20.2 ^b	23.7 ^a	23.1 ^a	25.1 ^a	0.50	0.04
♀(5)	22.3 ^a	23.3 ^a	21.0 ^a	24.3 ^a	0.48	0.44
♂♀(10)	21.2 ^b	23.5 ^a	22.0 ^a	24.7 ^a	0.41	0.10
Crude fiber (CF)						
♂(5)	45.6 ^b	56.4 ^a	54.5 ^a	56.7 ^a	1.26	0.04
♀(5)	51.5 ^a	54.5 ^a	49.4 ^a	55.6 ^a	1.13	0.35
♂♀(10)	48.5 ^b	55.4 ^a	51.9 ^a	56.1 ^a	1.01	0.04

a, b: Averages with the same letters on the same line are not significantly different at the 5% level; ESM: Standard Error on the Average; P: Probability; PMCA0: *P. maximum* + diet containing 0% of *A. glabrata*; PMCA10: *P. maximum* + diet containing 10% of *A. glabrata*; PMCA15: *P. maximum* + diet containing 15% of *A. glabrata*; PMCA20: *P. maximum* + diet containing 20% of *A. glabrata*.




Figure 1: Effect of *A. glabrata* levels in the diet on DM, OM, CP, and CF intake in guinea pigs fed with *P. maximum* according to sex
ab: Averages with the same letters on the same line are not significantly different at the 5%

Table 5: Apparent Digestibility Coefficients (ADC) of nutrients in guinea pigs fed with *P. maximum* associated with inclusion level of *A. glabrata*.

ADC (%) Sex	Treatments				ESM	P
	PMCA0	PMCA10	PMCA15	PMCA20		
ADC DM						
♂(5)	92.1 ^a	91.8 ^a	91.7 ^a	91.2 ^a	0.33	0.98
♀(5)	93.2 ^a	93.1 ^a	91.4 ^a	91.3 ^a	0.49	0.43
♂♀(10)	92.6 ^a	92.5 ^a	91.6 ^a	91.2 ^a	0.29	0.76
ADC OM						
♂(5)	92.3 ^a	92.0 ^a	92.0 ^a	91.5 ^a	0.32	0.92
♀(5)	93.3 ^a	93.2 ^a	91.6 ^a	91.5 ^a	0.47	0.45
♂♀(10)	92.8 ^a	92.6 ^a	91.8 ^a	91.5 ^a	0.28	0.76
ADC CP						
♂(5)	95.5 ^a	94.2 ^a	94.1 ^a	94.0 ^a	0.25	0.43
♀(5)	95.6 ^a	95.3 ^a	94.3 ^a	94.1 ^a	0.37	0.67
♂♀(10)	95.6 ^a	94.8 ^a	94.2 ^a	94.1 ^a	0.23	0.76
ADC CF						
♂(5)	90.8 ^a	90.3 ^a	90.2 ^a	90.2 ^a	0.42	0.90
♀(5)	91.4 ^a	91.1 ^a	91.0 ^a	90.8 ^a	0.60	0.34
♂♀(10)	91.1 ^a	90.7 ^a	90.6 ^a	90.5 ^a	0.37	0.65

ab: Averages with the same letters on the same line are not significantly different at the 5% level; ESM: Standard Error on the Average; P: Probability; ADC: apparent digestibility coefficient ; PMCA0: *P. maximum* + diet containing 0% of *A. glabrata*; PMCA10: *P. maximum* + diet containing 10% of *A. glabrata*; PMCA15: *P. maximum* + diet containing 15% of *A. glabrata*; PMCA20: *P. maximum* + diet containing 20% of *A. glabrata*.

the bacterial flora of guinea pigs fed with *P. maximum*

The rate of Enterobacteria and Lactobacilli in guinea pigs fed *P. maximum* did not significantly ($p > 0.05$) change with the inclusion level of *A. glabrata* in the diet (Table 6). The highest rate of Enterobacteria (4.3 CFU/ ml) and Lactobacilli (4.6 CFU/ ml) was obtained with the 20% *A. glabrata* diet. Enterobacteria levels were comparable ($p > 0.05$) to those of Lactobacilli in guinea pigs fed with *P. maximum* regardless of diet (Figure 3). Comparison of the levels of these bacteria between males and females revealed no significant difference (Figure 4).

DISCUSSION

Effect of inclusion level of *Arachis glabrata* in diet, on ingestion of *Panicum maximum* in guinea pigs

Compared with the control diet, *P. maximum* ingestion (160.3 g

DM/ animal/ d) was significantly higher with *A. glabrata*-containing diets. This shows that the protein intake of this legume improved the palatability of guinea pigs. Moreover, several works by Kouakou




Figure 2: Effect of sex on apparent digestibility coefficient of DM, OM, CP and CF in guinea pigs fed with different levels of *A. glabrata*. ADC: apparent digestibility coefficient; PMCA0: *P. maximum* + diet containing 0% of *A. glabrata*; PMCA10: *P. maximum* + diet containing 10% of *A. glabrata*; PMCA15: *P. maximum* + diet containing 15% of *A. glabrata*; PMCA20: *P. maximum* + ration containing 20% of *A. glabrata*.

Table 6: Effect of inclusion level of *Arachis glabrata* in diet on the variation of Enterobacteria and Lactobacilli levels in guinea pigs fed *P. maximum*.

Genus bacterial sex	Treatments				ESM	P
	PMCA0	PMCA10	PMCA15	PMCA20		
LOG10 (Enterobacteria) (CFU/ml)						
♂(5)	3.5 ^a	4.3 ^a	4.3 ^a	4.5 ^a	0.09	0.05
♀(5)	4.0 ^a	3.8 ^a	3.9 ^a	4.2 ^a	0.07	0.13
♂♀(10)	3.7 ^a	4.0 ^a	4.1 ^a	4.3 ^a	0.08	0.73
LOG10 (Lactobacillii) (CFU/ml)						
♂(5)	4.4 ^a	4.8 ^a	4.9 ^a	5.0 ^a	0.09	0.20
♀(5)	4.1 ^a	4.2 ^a	3.9 ^a	4.3 ^a	0.06	0.68
♂♀(10)	4.2 ^a	4.5 ^a	4.4 ^a	4.6 ^a	0.10	0.46

a, b: Averages with the same letters on the same line are not significantly different at the 5% level; ESM: Standard Error on the Average; P: Probability; PMCA0: *P. maximum* + diet containing 0% of *A. glabrata*; PMCA10: *P. maximum* + diet containing 10% of *A. glabrata*; PMCA15: *P. maximum* + diet containing 15% of *A. glabrata*; PMCA20: *P. maximum* + diet containing 20% of *A. glabrata*.




Figure 3: Effect of the inclusion level of *Arachis glabrata* in the diet on the variation of the level of Enterobacteria and Lactobacilli in guinea pigs fed with *P. maximum*. *P. maximum* + ration containing 0% of *A. glabrata*; PMCA10: *P. maximum* + diet containing 10% of *A. glabrata*; PMCA15: *P. maximum* + diet containing 15% of *A. glabrata*; PMCA20: *P. maximum* + diet containing 20% of *A. glabrata*.




Figure 4: Effect of sex on apparent digestibility coefficient of DM, OM, CP and CF in guinea pigs fed with different levels of *A. glabrata*. ADC: apparent digestibility coefficient; PMCA0: *P. maximum* + diet containing 0% of *A. glabrata*; PMCA10: *P. maximum* + diet containing 10% of *A. glabrata*; PMCA15: *P. maximum* + diet containing 15% of *A. glabrata*; PMCA20: *P. maximum* + ration containing 20% of *A. glabrata*.

et al. [10]; Miegoue et al. [8] and Miegoue et al. [9] illustrate this observation.

The best intake of these grasses obtained with the ration containing 20% of *A. glabrata* is explained by the fact that protein supplements of plant origin (more adapted to the herbivore diet of guinea pigs) would promote a sufficient proliferation of intestinal microorganisms involved in digestion in herbivores. This would favor the increase of food fermentation and digestive transit which would thus de-clutter the caecum with the consequent increase in food intake (Kouakou et al. [10], Ramirez-Riviera et al. [16], Dougnon et al. [17]).

The low intake of the compound feed in rations containing the legume compared to the control diet could be explained by the substitution phenomenon illustrated [18]. In fact, when a food supplementing self-service feed is distributed separately, a portion of the digestive tract of the animal which would mean that the presence of *A. glabrata* in the diet increased the palatability of *P. maximum* in the animal that has more ingested the grass at the expense of the compound feed. Moreover the inclusion of *A. glabrata* in the diet would have increased its capacity for congestion and satiety in the guinea pig, thus reducing ingestion of rations containing this legume contrary to the control diet. The level of intake of total Dry Matter (DM) and Organic Matter (OM) from *A. glabrata*-containing diets higher in male guinea pigs compared females could be explained by the fact that in general in adulthood, males have a high weight compared to that of females. As a result, they tend to ingest more food because food intake is very often correlated with the weight of the animal. Indeed according to many authors, dietary intake in male guinea pigs is higher than that in females Noubissi et al. [6], Miegoue et al. [8].

Effect of inclusion level of *Arachis glabrata* in the diet on the digestibility of *Panicum maximum* in guinea pigs

The lack of a significant difference between the Apparent Digestibility Coefficients (ADC) of nutrients from different diets in guinea pigs fed *P. maximum* could be explained by the fact that the diets used in this trial were iso nitrogenated and of little variable chemical composition. Indeed, according [18], diet is the factor that has the strongest influence on digestibility [19]. Also reported that



one of the environmental factors that seems most likely to modify the bacterial colonization process of the gastrointestinal tract is the food ingested by the host. Digestibility will therefore be affected by the quality and physical state of the ration, that is to say the form in which the food is presented to the animal as much as its chemical composition. These factors condition the action of the microbial flora and digestive juices. The rations served to guinea pigs were in powder form, which favored both ingestion and digestive utilization of the nutrients.

Effect of inclusion level of *Arachis glabrata* in rations on the caecal rate of *Enterobacteria* or *Lactobacillus* bacterial flora in guinea pigs fed *P. maximum*

The level of inclusion of *Arachis glabrata* in diet had no significant effect on the variation of bacterial flora in the enteric bacterial and lactobacilli species in guinea pigs. This shows a certain balance between this bacterial population. Indeed, According Losson [20] the balance of this flora is unstable, easily altered by abrupt food change or the administration of a narrow-spectrum antibiotic Gram (+), which can lead to enteropathies. The absence of variation of the flora would therefore mean that the inclusion of *Arachis glabrata* in the experimental rations did not alter the composition of the flora and consequently the digestive use of the nutrients. Similar results were obtained by Miegoue et al. [8] who had shown that the inclusion of legumes in the guinea pig diet did not significantly affect the caecal flora. Diets thus composed were therefore adapted to guinea pig feeding. Compared with the rate of *Enterobacteria* (Gram -), *Lactobacilli* (Gram +) were the most numerous, whatever the diet used. This result reflects the quality of the experimental foods. In fact, according to Andreu and Lormeau most of the caecal microbial population is predominantly composed of Gram (+) and anaerobic organisms, Gram (-) being present in a smaller quantity [21-23].

CONCLUSION

Evaluation of the effect of different dietary supplementation levels with *Arachis glabrata* on the *in vivo* digestibility of *Panicum maximum* in guinea pigs showed that:

- The ration containing 20% of *Arachis glabrata* significantly improved the intake of *Panicum maximum*. Males had better ingested DM and OM of diet containing legumes than females.
- Inclusion levels of *Arachis glabrata* in the diet did not improve digestive utilization of nutrients ingested by guinea pigs.
- Cecal flora was not significantly affected by the inclusion level of *Arachis glabrata* when guinea pigs were fed with *P. maximum*.

Thus, *Arachis glabrata* can be used up to 20% in cavy diet without any in food intake or utilization.

REFERENCES

1. Niba AT, Meutchieye F, Fon D, Laisin AG, Taboh H, Njakoi H, et al. Current situation of cavy production in Cameroon: Challenges and opportunities. *Livestock Research for Rural Development*. 2012 ; 24. <https://goo.gl/IVk5Tw>
2. FAO. FAO role in nutrition. 2017.
3. FAO. Livestock farming and food security. 2011.
4. Noubissi M, Tendongkeng F, Zougou T, Miegoue E, Lemoufouet J, Boukila B, et al. Effect of the complementation with *Tithonia diversifolia* on the evolution of post-partum weight and pre-weaning growth of cavy (Cavia porcellus L.). *Livestock Research for Rural Development*. 2013; 25. <https://goo.gl/tRa4Nj>
5. Tendongkeng F, Boukila B, Pamo TE, Mboko AV, Zogang FB, et Matumuini NEF. 2011. Direct and residual Effects of different level of Nitrogen fertilisation on the chemical composition of *Brachiaria ruziziensis* at the flowering stage in the west Cameroon. *International Journal of Biological and Chemical Sciences*. 2011; 5: 570-585. <https://goo.gl/dPHYCL>
6. Noubissi MNB, Tendongkeng F, Zougou TG, et Pamo TE. Effect of different supplementation level of *Tithonia diversifolia* Hemsl. Leaves on the ingestion and the *in vivo* digestibility of *Penisetum purpureum* K. Schum. On cavy (Cavia porcellus L.). *Tropicicultura*. 2014 ; 3: 138-146.
7. Bindelle J, et Picron P. The cavy : a small herbivorous animal easy to feed in small plots. *Troupeaux et Cultures des Tropiques : Special rodents breeding*, Kinshasa. RDC. CAVTK. 2013; 1-10.
8. Miegoue E, Tendongkeng F, Lemoufouet J, Mweugang Nguoupo N, Noubissi M, Fogang MD, et al. intake and digestibility of *Penisetum purpureum* associate to one legume (*Arachis glabrata*, *Calliandra calothyrsus* ou *Desmodium intortum*) as protein source on cavy. *Livestock Research for Rural Development*. 2016a; 28. <https://goo.gl/e76n7E>
9. Miegoue E, Tendongkeng F, Lemoufouet J, Noubissi M N B, Mweugang N N, Zougou TG, et al. pre-weaning growth of cavy fed on *Panicum maximum* supplemented with diet containing *Arachis glabrata*, *Calliandra calothyrsus* or *Desmodium intortum*. *Int J Biol Chem Sci*. 2016b; 10: 313-325. <https://goo.gl/PpsTts>
10. Kouakou N'GDV, Thys E, Assidjo EN, et Grongnet JF. Ingestion and *in vivo* digestibility of *Panicum maximum* associated to three complements: *Jatropha curcas* seed cake, tourteau de coton seed cake (*Gossypium hirsutum*) and *Euphorbia heterophylla* on cavy (Cavia porcellus L.). *Tropicicultura*. 2010; 28: 173-177. <https://goo.gl/xtUkub>
11. Michelland R, Combes S, Monteils V, Bayourthe C, Cauquil L, Enjalbert F, et al. compare Analysis of digestive ecosystems of rumen of caw and rabbits caecum. *Prod Anim*. 2012 ; 25 : 395-406. <https://goo.gl/76n8pf>
12. Pamo TE, Niba A T, Fonteh FA, Tendongkeng F, Kana JR, Boukila B, et al. Effect of the supplementation with *Moringa oleifera* or with multivitamin blocks on the evolution of post-partum weight and pre-weaning growth of cavy (Cavia porcellus L.). *Livestock Research for Rural Development*. 2005; 17. <https://goo.gl/fJhSp7>
13. AOAC. (Association of Official Analytical Chemist). 15th edition. Washington DC. Official method of analysis. 1990; 10.
14. Benson HJ. Microbiological applications. Laboratory manual in general microbiology. Complete version. Eighth edition. Mc. Graw Hill higher Education International edition ISBN 0-07-112169-2. 2002; 478.
15. Steele R G, Torrie JH. Principles and procedures of statistics. Mc Graws Hill Book C. 1980; 633.
16. Ramirez-Rivera U, Sangines-Garcia JR, Escobedo-Mex JG, Cen-Chuc F, Rivera-Lorca JA, Lara-Lara PE. Effect of diet inclusion of *Tithonia diversifolia* on feed intake, digestibility and nitrogen balance in tropical sheep. *Agroforest System*. 2010 ; 80: 295-302. <https://goo.gl/pTvkSg>
17. Dougnon TJ, Aboh BA, Kpodekon TM. Ingestion, digestibility and ponderal growth on cricetomies (*Cricetomys gambianus*) fed on *Moringa oleifera* and *Leucaena leucocephala* based concentrate. *Livestock Research for Rural Development*. 2011; 23: 15-27. <https://goo.gl/roFGhW>
18. Rivière R. Domestic ruminants feeding Manual in tropical milieu. m a n u a l and precise of breeding Collection. Ministry of cooperation and development. 1991 ; 529.
19. Rainbaud P, et Ducluzeau R. Study of bacterian colonization bactérienne gastro-intestinal tract using experimental model. *Rev. Sc. Tech. off. Int. Epiz*. 1989; 8: 361-373. <https://goo.gl/7wR9ky>
20. Losson B. The cavy and his principales diseases. herd and Cultures of Tropics : Special rodents breeding, Kinshasa, RDC. Laboratory of Parasitology, Faculty of veterinary Medicine, University of Liège, Bd de Colonster 20 Bât. B43, 4000 Liège, Belgique. Thesis. Ecole Nationale Vétérinaire d'Alfort, Alfort. 2013; 122.



21. Andreu de Lapierre E. The cavy. Medecine pratique Dictionary of NCA, Ed. MED'COM. 2001; 18-29.
22. Lormeau E. Contribution of the study of *Cavia porcellus* (Linne, 1758): Atlas of radiographies and osteology. Exercice Theisis, veterinary Medicine, Toulouse. 2010; 207.
23. Pamo T E, Niba A T, Fonteh FA, Tendonkeng F, Kana JR, Boukila B, et al. Effect of the supplementation with *Moringa oleifera* or with multivitaminized blocks on the evolution of post-partum weight and pre-weaning growth of cavy (Cavia porcellus L.). Livestock Research for Rural Development. 2005; 17. <https://goo.gl/6fqXEE>